



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

that in order to exhibit this properly, the section must be exceedingly thin. He stated that an optical examination had proved that the direction of the main lines of the markings corresponded with or was at right angles to that of the crystallographic axes of the Muscovite. He exhibited a plate of the mica shown distinctly to be a twin by the two different groupings of Magnetite markings. Examination in the polariscope confirmed this structure. Thus, frequently, the crystalline structure of the mica and the direction of its axes may be ascertained by an inspection of these markings alone. It appeared, therefore, that the form and direction of the markings was determined, not by independent crystallization of the Magnetite forming them, but in part at least by the Muscovite from which it had probably been derived. These markings are, in some respects, pseudomorphs after Muscovite. He thought that the statement in Dana's Mineralogy (p. 150), referring to these markings, that "the branching at angles of  $60^{\circ}$  indicates composition parallel to a dodecahedral face," was misleading, implying that this form was produced by an inherent property of the Magnetite, and not, as he thought now appears, by the crystalline structure of the Muscovite.

SEPTEMBER 24, 1877.

*A New Locality for Asbolite.*—Mr. LEWIS stated that he had found Asbolite at Flourtown, Montgomery Co., a new locality for this mineral. It is found in iron ore mines as an incrustation upon Psilomelane. It is of a bluish-black color, is as soft as graphite, and gives a shining streak when scratched by the nail. The blowpipe indicates a considerable percentage of cobalt.

*A New Locality for Fluorite.*—Mr. W. W. JEFFERIS stated that a few days since he was shown a massive specimen of Fluor-spar of a deep purple color, which was found in the limestone near the village of Howellville, in Tredyffrin Township, Chester County, Pa. This is the third locality of fluor in this county.

*Epidote in Molybdenite.*—Mr. LEWIS mentioned that while examining some Molybdenite from Frankford, Phila., he had found plates of a transparent hard mineral, of a light greenish-yellow color, somewhat resembling Wulfenite, occurring in thin layers and minute scales between the foliæ of the Molybdenite, and sometimes coating it as a thin film. It was not until after a careful examination that it was proved to be Epidote in an unusual form and situation.

OCTOBER 22, 1877.

*A New Locality for Millerite.*—Mr. THEO. D. RAND announced the discovery of Millerite in Dolomite, from the Soapstone quarry on the Schuylkill, in Philadelphia, near the Montgomery County line. It occurred in capillary crystals in cavities of the Dolomite.